



Three Rivers

Sexual and Reproductive Health Profile

The Sexual and Reproductive Health Profiles have been developed by Public Health England to support local authorities and public health leads monitor the sexual and reproductive health of their population and the performance of local public health related systems.

The profiles provide a snapshot of sexual and reproductive health across a range of topics including teenage pregnancy, abortions, contraception, HIV and sexually transmitted infections. Wider influences on sexual health such as alcohol use, and other topics relating to teenage pregnancy such as education and deprivation level, are also included.

This output report provides selected data presentations showing how your local authority compares to the England value and other local authorities in your region.

Please refer to the online tool for accompanying advice and further presentations including interactive maps and other options for grouping and comparing your local authority.

Contents

Spine Charts

Key Indicators	Page 2
HIV & STI	Page 3
Reproductive Health	Page 3
Teenage Pregnancy	Page 3
Wider Determinants of Health	Page 3

Trends

Key Indicators	Page 5
HIV & STI	Page 7
Reproductive Health	Page 9
Teenage Pregnancy	Page 10
Wider Determinants of Health	Page 11

Tartan Rugs

Key Indicators	Page 12
HIV & STI	Page 12
Reproductive Health	Page 13
Teenage Pregnancy	Page 13
Wider Determinants of Health	Page 14

Bar Charts

Key Indicators	Page 15
HIV & STI	Page 17
Reproductive Health	Page 19
Teenage Pregnancy	Page 20
Wider Determinants of Health	Page 21

Definitions

Page 22



© Crown Copyright and database rights 2015, Ordnance Survey 100016969
OpenStreetMap contributors ODbL

<http://fingertips.phe.org.uk/profile/sexualhealth>

Produced by Public Health England.

For enquiries please contact profilefeedback@phe.gov.uk


Three Rivers

Spine Charts

Key

Significance compared to goal / England average:

- Significantly worse
- Significantly lower
- Not significantly different
- Significantly higher
- Significantly better
- Significance not tested



Key Indicators

	Period	Local count	Local value	Eng. value	Eng. worst / lowest	Range	Eng. best / highest
Syphilis diagnosis rate / 100,000	2013	1	1.1	5.9	90.9		0.0
Gonorrhoea diagnosis rate / 100,000	2013	22	24.8	52.9	533.2		1.3
Chlamydia detection rate / 100,000 aged 15-24 (PHOF indicator 3.02)	2013	117	1199	2016	643		5758
< 1900 1900 to 2300 ≥ 2300							
Chlamydia proportion aged 15-24 screened	2013	1,451	14.9	24.9	8.5		100
All new STI diagnoses (exc Chlamydia aged <25) / 100,000	2013	487	858	832	3269		172
HIV testing uptake, MSM (%)	2013	88	94.6	94.8	82.5		100
HIV late diagnosis (%) (PHOF indicator 3.04)	2011 - 13	8	66.7	45.0	100.0		14.3
< 25 25 to 50 ≥ 50							
HIV diagnosed prevalence rate / 1,000 aged 15-59	2013	57	1.09	2.14	0.24		14.70
Under 18s conception rate / 1,000 (PHOF indicator 2.04)	2013	35	20.2	24.3	44.0		7.6
Under 18s conceptions leading to abortion (%)	2013	*	- *	51.1	31.4		89.5
Sexual offences rate / 1,000 (PHOF indicator 1.12iii)	2013/14	32	0.36	1.01	0.35		2.43

Note: * - Disclosure control applied

HIV & STI

	Period	Local count	Local value	Eng. value	Eng. worst / lowest	Range	Eng. best / highest
Syphilis diagnosis rate / 100,000	2013	1	1.1	5.9	90.9		0.0
Gonorrhoea diagnosis rate / 100,000	2013	22	24.8	52.9	533.2		1.3
Chlamydia detection rate / 100,000 aged 15-24 (PHOF indicator 3.02)	2013	117	1199	2016	643		5758
<div style="display: flex; justify-content: space-between; width: 100%; font-size: small;"> < 1900 1900 to 2300 ≥ 2300 </div>							
Chlamydia proportion aged 15-24 screened	2013	1,451	14.9	24.9	8.5		100
Genital warts diagnosis rate / 100,000	2013	98	110.4	133.4	288.6		30.0
Genital herpes diagnosis rate / 100,000	2013	52	58.6	58.8	182.9		10.8
All new STI diagnoses (exc Chlamydia aged <25) / 100,000	2013	487	858	832	3269		172
STI testing rate (exc Chlamydia aged < 25) / 100,000	2013	7,449	13131	14685	4171		53921
STI testing positivity (exc Chlamydia aged <25) %	2013	487	6.5	5.7	3.7		11.2
HIV testing uptake, MSM (%)	2013	88	94.6	94.8	82.5		100
HIV testing uptake, women (%)	2013	901	79.2	75.8	29.0		94.4
HIV testing uptake, men (%)	2013	748	86.3	84.9	58.4		95.9
HIV testing coverage, MSM (%)	2013	68	85.0	86.1	63.2		100
HIV testing coverage, women (%)	2013	812	60.8	65.6	26.0		87.7
HIV testing coverage, men (%)	2013	688	78.3	77.5	44.5		89.4
HIV late diagnosis (%) (PHOF indicator 3.04)	2011 - 13	8	66.7	45.0	100.0		14.3
<div style="display: flex; justify-content: space-between; width: 100%; font-size: small;"> < 25 25 to 50 ≥ 50 </div>							
HIV diagnosed prevalence rate / 1,000 aged 15-59	2013	57	1.09	2.14	0.24		14.70

Reproductive Health

	Period	Local count	Local value	Eng. value	Eng. worst / lowest	Range	Eng. best / highest
Pelvic inflammatory disease (PID) admissions rate / 100,000	2013/14	31	184.3	235.7	686.7		54.4
Ectopic pregnancy admissions rate / 100,000	2013/14	12	71.3	89.6	200.9		22.4
Cervical cancer registrations rate / 100,000	2010 - 12		7.2	9.2	18.2		3.4

Teenage Pregnancy


	Period	Local count	Local value	Eng. value	Eng. worst / lowest	Range	Eng. best / highest
Under 18s conception rate / 1,000 (PHOF indicator 2.04)	2013	35	20.2	24.3	44.0		7.6
Under 18s conceptions leading to abortion (%)	2013	*	- *	51.1	31.4		89.5
Under 18s births rate / 1,000	2013	6	3.5	7.8	20.1		0.0

Wider Determinants of Health

	Period	Local count	Local value	Eng. value	Eng. worst / lowest	Range	Eng. best / highest
Under 18s alcohol-specific hospital admissions rate / 100,000	2011/12 - 13/14	20	31.5	40.1	105.8		11.2
Percentage people living in 20% most deprived areas in England	2013	1,661	1.9	20.4	83.8		0.0
Under 16s in poverty (%) (PHOF indicator 1.01ii)	2012	1,905	11.0	19.2	37.9		5.8
GCSE achieved 5A*-C inc. Eng & Maths (%)	2012/13	763	76.9	60.8	38.1		81.9

Note: * - Disclosure control applied


Three Rivers

Wider Determinants of Health continued	Period	Local count	Local value	Eng. value	Eng. worst / lowest	Range	Eng. best / highest
Pupil absence (%) (PHOF indicator 1.03)	2012/13	184,702	4.78	5.26	6.31		4.10


Trends

Key Indicators


Syphilis diagnosis rate / 100,000




Gonorrhoea diagnosis rate / 100,000




Chlamydia detection rate / 100,000 aged 15-24 (PHOF indicator 3.02)




Chlamydia proportion aged 15-24 screened




All new STI diagnoses (exc Chlamydia aged <25) / 100,000




HIV testing uptake, MSM (%)




HIV late diagnosis (%) (PHOF indicator 3.04)




HIV diagnosed prevalence rate / 1,000 aged 15-59



Under 18s conception rate / 1,000 (PHOF indicator 2.04)




Under 18s conceptions leading to abortion (%)



Three Rivers


Key Indicators continued

Sexual offences rate / 1,000 (PHOF indicator 1.12iii)




HIV & STI


Syphilis diagnosis rate / 100,000




Gonorrhoea diagnosis rate / 100,000




Chlamydia detection rate / 100,000 aged 15-24 (PHOF indicator 3.02)




Chlamydia proportion aged 15-24 screened




Genital warts diagnosis rate / 100,000




Genital herpes diagnosis rate / 100,000




All new STI diagnoses (exc Chlamydia aged <25) / 100,000




STI testing rate (exc Chlamydia aged < 25) / 100,000



STI testing positivity (exc Chlamydia aged <25) %




HIV testing uptake, MSM (%)




Three Rivers

HIV & STI continued


HIV testing uptake, women (%)




HIV testing uptake, men (%)




HIV testing coverage, MSM (%)




HIV testing coverage, women (%)




HIV testing coverage, men (%)



HIV late diagnosis (%) (PHOF indicator 3.04)




HIV diagnosed prevalence rate / 1,000 aged 15-59




Reproductive Health


Pelvic inflammatory disease (PID) admissions rate / 100,000



Ectopic pregnancy admissions rate / 100,000




Cervical cancer registrations rate / 100,000




Three Rivers

Teenage Pregnancy


Under 18s conception rate / 1,000 (PHOF indicator 2.04)



Under 18s conceptions leading to abortion (%)




Under 18s births rate / 1,000




Wider Determinants of Health


Under 18s alcohol-specific hospital admissions rate / 100,000




Percentage people living in 20% most deprived areas in England




Under 16s in poverty (%) (PHOF indicator 1.01ii)



GCSE achieved 5A*-C inc. Eng & Maths (%)



Pupil absence (%) (PHOF indicator 1.03)



Three Rivers

Tartan Rugs

Key Indicators

Indicator	Period	England	Barbergh	Basildon	Bedford	Braintree	Breckland	Brentwood	Broadland	Broxbourne	Cambridge	Castle Pt	Central Beds	Chelmsford	Colchester	Dacorum	E Cambs	East Herts	Epping For	Fenland	Forest Heath	Gt Yarmouth	Harlow	Hertsme	Hunts	Ipswich	KL & W Norf	Luton	Maldon	Mid Suffolk	North Herts	N Norfolk	Norwich	Peterboro	Rochford	S Cambs	S Norfolk	Southend	St Albans	St Edmundsbury	Stevenage	Suff Coastal	Tendring	Thr Rivers	Thurrock	Uttlesford	Wafford	Waveney	Welw Hat'd
Syphilis diagnosis rate / 100,000	2013	5.9	0.0	2.3	5.7	2.0	2.3	0.0	3.2	2.1	3.2	0.0	1.9	4.1	3.4	1.4	1.2	2.9	3.2	1.0	1.6	5.1	4.8	2.0	0.6	3.0	0.7	2.9	1.6	2.0	2.3	2.0	7.4	2.1	0.0	0.7	0.0	4.6	1.4	2.7	7.1	0.8	2.9	1.1	2.5	0.0	6.5	0.9	1.8
Gonorrhoea diagnosis rate / 100,000	2013	52.9	8.0	23.2	62.2	14.2	14.4	12.2	14.4	27.5	41.5	24.9	18.1	19.5	23.3	12.3	7.1	15.8	38.9	9.4	14.8	11.3	33.9	35.7	12.3	31.2	8.7	50.5	24.2	16.4	22.6	5.9	49.9	42.4	19.1	11.3	8.7	43.5	29.6	8.1	40.1	11.3	13.0	24.8	31.3	13.5	42.5	7.8	35.7
Chlamydia detection rate / 100,000 aged 15-24 (PHOF indicator 3.02)	2013	2016	931	1556	2993	1642	1123	1283	1222	1397	1484	1748	1190	1476	2226	4232	1467	1519	1753	1790	911	1187	2687	979	1780	2861	1379	1799	1631	945	1598	1185	1756	2489	1556	1105	1508	2299	948	1121	4931	835	1607	1199	1529	1372	1393	2026	913
			< 1900	1900 to 2300	> 2300																																												
Chlamydia proportion aged 15-24 screened	2013	24.9	17.8	17.0	28.0	18.7	15.2	17.0	16.8	12.8	30.3	21.3	16.0	20.8	35.3	66.9	25.3	15.4	21.1	28.9	12.1	15.3	27.8	12.4	32.9	43.0	15.4	19.7	18.8	13.7	14.1	15.1	24.1	22.7	22.1	20.6	16.0	28.1	16.6	17.6	37.0	15.4	24.5	14.9	18.0	18.2	16.8	21.6	11.1
All new STI diagnoses (exc Chlamydia aged <25) / 100,000	2013	832	520	614	901	531	565	555	579	689	775	549	530	715	873	713	362	669	826	392	420	741	1155	834	561	887	792	885	439	505	487	412	1193	798	481	481	570	721	809	508	980	571	580	858	667	572	1294	667	747
HIV testing uptake, MSM (%)	2013	94.8	96.0	97.3	91.4	96.9	100	98.4	99.1	90.3	97.7	94.4	95.6	97.1	92.4	95.0	100	91.7	93.5	96.0	97.7	94.0	90.4	90.7	94.0	91.1	87.8	93.4	96.4	92.0	93.0	97.0	97.4	97.1	92.9	98.1	92.7	94.5	94.5	97.4	93.2	92.7	83.0	94.6	95.4	93.2	96.3	94.9	92.1
HIV late diagnosis (%) (PHOF indicator 3.04)	2011 - 13	45.0	57.1	72.7	77.3	66.7	54.5	100.0	42.9	53.3	40.5	50.0	46.7	46.4	47.1	80.0	-	50.0	42.9	-	45.5	38.4	42.3	47.1	54.5	40.0	43.8	50.7	-	33.3	35.7	70.0	47.2	61.9	46.2	50.0	63.6	49.0	53.3	46.2	50.0	40.0	30.0	66.7	63.2	-	51.9	62.5	50.0
			< 25	25 to 50	> 50																																												
HIV diagnosed prevalence rate / 1,000 aged 15-59	2013	2.14	0.73	1.47	2.17	0.83	0.87	1.43	0.62	1.31	1.43	0.54	1.29	1.35	1.26	1.21	0.60	0.96	1.21	0.96	1.08	1.07	2.67	1.86	1.18	1.38	0.84	4.07	0.62	0.75	1.26	0.92	2.17	1.72	0.73	0.80	0.82	2.99	1.05	0.86	2.33	0.51	0.66	1.09	2.10	0.66	2.79	0.96	1.95
Under 18s conception rate / 1,000 (PHOF indicator 2.04)	2013	24.3	15.4	32.9	22.0	22.3	19.3	12.8	15.5	12.5	21.9	17.4	19.9	15.8	20.6	21.0	8.6	12.2	15.4	23.1	19.9	26.5	38.3	17.0	15.4	29.4	28.7	24.4	27.1	16.3	15.5	19.0	31.4	33.4	16.8	13.4	16.0	26.6	7.6	19.1	29.1	12.8	30.3	20.2	36.1	13.3	18.5	23.3	16.7
Under 18s conceptions leading to abortion (%)	2013	51.1	40.0	51.4	50.7	55.0	55.6	89.5	54.3	77.3	42.9	44.4	58.7	57.8	44.1	56.9	-	64.7	78.8	-	-	37.5	50.9	53.1	54.2	43.9	36.1	52.6	72.4	-	52.8	46.9	42.9	46.6	63.0	59.5	54.1	37.3	-	47.2	50.0	60.0	40.8	-	57.4	59.1	80.0	31.9	74.2
Sexual offences rate / 1,000 (PHOF indicator 1.12iii)	2013/14	1.01	0.74	1.13	0.82	0.69	1.14	0.53	0.68	0.55	0.86	0.76	0.55	0.89	1.32	0.71	0.58	0.48	0.59	0.85	0.76	1.20	1.20	0.48	0.76	1.29	0.98	0.91	0.48	0.76	0.48	1.03	1.66	1.45	0.51	0.77	0.53	1.41	0.56	0.76	1.30	0.60	1.16	0.36	0.99	0.52	0.97	1.07	0.76

HIV & STI

Syphilis diagnosis rate / 100,000	2013	5.9	0.0	2.3	5.7	2.0	2.3	0.0	3.2	2.1	3.2	0.0	1.9	4.1	3.4	1.4	1.2	2.9	3.2	1.0	1.6	5.1	4.8	2.0	0.6	3.0	0.7	2.9	1.6	2.0	2.3	2.0	7.4	2.1	0.0	0.7	0.0	4.6	1.4	2.7	7.1	0.8	2.9	1.1	2.5	0.0	6.5	0.9	1.8	
Gonorrhoea diagnosis rate / 100,000	2013	52.9	8.0	23.2	62.2	14.2	14.4	12.2	14.4	27.5	41.5	24.9	18.1	19.5	23.3	12.3	7.1	15.8	38.9	9.4	14.8	11.3	33.9	35.7	12.3	31.2	8.7	50.5	24.2	16.4	22.6	5.9	49.9	42.4	19.1	11.3	8.7	43.5	29.6	8.1	40.1	11.3	13.0	24.8	31.3	13.5	42.5	7.8	35.7	
Chlamydia detection rate / 100,000 aged 15-24 (PHOF indicator 3.02)	2013	2016	931	1556	2993	1642	1123	1283	1222	1397	1484	1748	1190	1476	2226	4232	1467	1519	1753	1790	911	1187	2687	979	1780	2861	1379	1799	1631	945	1598	1185	1756	2489	1556	1105	1508	2299	948	1121	4931	835	1607	1199	1529	1372	1393	2026	913	
			< 1900	1900 to 2300	> 2300																																													

Comparison with respect to England value / goal



Note: * - Disclosure control applied

<http://fingertips.phe.org.uk/profile/sexualhealth>

E07000102 Three Rivers

© Crown Copyright 2015. Updated 2 June 2015

HIV & STI continued

Period	England	Babergh	Basildon	Bedford	Braintree	Breckland	Brentwood	Broadland	Broxbourne	Cambridge	Castle Pt	Central Beds	Chelmsford	Colchester	Dacorum	E Cambs	East Herts	Epping For	Fenland	Forest Heath	Gt Yarmouth	Harlow	Hertsmere	Hunts	Ipswich	KL & W Norf	Luton	Maldon	Mid Suffolk	North Herts	N Norfolk	Norwich	Peterboro	Rochford	S Cambs	S Norfolk	Southend	St Albans	St Edm'bury	Stevenage	Suff Coastal	Tending	Thr Rivers	Thurrock	Uttlesford	Watford	Waveney	Welw Hat'd	
Chlamydia proportion aged 15-24 screened	2013	24.9	17.8	17.0	28.0	18.7	15.2	17.0	16.8	12.8	30.3	21.3	16.0	20.8	35.3	66.9	25.3	15.4	21.1	28.9	12.1	15.3	27.8	12.4	32.9	43.0	15.4	19.7	18.8	13.7	14.1	15.1	24.1	22.7	22.1	20.6	16.0	28.1	16.6	17.6	67.0	15.4	24.5	14.9	18.0	18.2	16.8	21.6	11.1
Genital warts diagnosis rate / 100,000	2013	133.4	97.8	133.2	136.3	129.4	97.1	122.9	99.0	125.9	137.4	127.0	109.6	147.6	188.6	123.4	69.3	128.4	132.5	67.7	84.0	126.1	156.0	123.1	125.1	131.6	109.7	127.3	96.9	92.2	76.3	54.0	190.7	130.4	133.5	101.3	82.6	148.7	126.1	119.2	120.3	105.4	140.3	110.4	131.6	115.7	164.6	103.7	124.9
Genital herpes diagnosis rate / 100,000	2013	58.8	43.2	57.8	70.3	48.5	46.3	59.4	43.1	61.4	44.7	41.9	44.2	69.7	75.0	39.5	17.6	62.4	64.2	28.1	41.2	75.8	79.8	64.5	40.3	77.3	60.6	48.1	54.9	48.2	38.9	28.5	100.5	47.2	34.6	33.1	50.0	60.1	56.4	52.0	70.8	40.2	41.9	58.6	60.8	45.5	75.2	61.4	42.8
All new STI diagnoses (exc Chlamydia aged <25) / 100,000	2013	832	520	614	901	531	565	555	579	689	775	549	530	715	873	713	362	669	826	392	420	741	1155	834	561	887	792	885	439	505	487	412	1193	798	481	481	570	721	809	508	980	571	580	858	667	572	1294	667	747
STI testing rate (exc Chlamydia aged < 25) / 100,000	2013	14685	8223	10722	14134	9542	8114	10500	9389	11694	13722	8253	9632	14734	15041	11049	6779	11685	12964	6839	8596	9732	15228	12968	9717	13597	11153	15167	8038	7675	10916	6896	17210	14451	8063	8281	9116	12329	14561	9930	20220	8615	9256	13131	12583	9159	19238	8593	13188
STI testing positivity (exc Chlamydia aged <25) %	2013	5.7	6.3	5.7	6.4	5.6	7.0	5.3	6.2	5.9	5.6	6.6	5.5	4.9	5.8	6.5	5.3	5.7	6.4	5.7	4.9	7.6	7.6	6.4	5.8	6.5	7.1	5.8	5.5	6.6	4.5	6.0	6.9	5.5	6.0	5.8	6.3	5.8	5.5	5.1	4.8	6.6	6.3	6.5	5.3	6.2	6.7	7.8	5.7
HIV testing uptake, MSM (%)	2013	94.8	96.0	97.8	91.4	96.9	100	98.4	99.1	90.3	97.7	94.4	95.6	97.1	92.4	95.0	100	91.7	93.5	96.0	97.7	94.0	90.4	90.7	94.0	91.1	87.8	93.4	96.4	92.0	93.0	97.0	97.4	97.1	92.9	98.1	92.7	94.5	94.5	97.4	93.2	92.7	88.0	94.6	95.4	93.2	96.3	94.9	92.1
HIV testing uptake, women (%)	2013	75.8	78.6	77.9	73.3	90.0	78.1	82.8	78.2	76.6	82.3	70.5	70.7	81.0	77.2	84.4	71.2	82.5	77.4	87.4	73.3	76.3	81.9	77.7	64.6	63.3	72.4	87.5	70.2	73.0	77.0	80.7	80.9	75.7	80.5	78.9	76.6	75.1	85.2	68.9	59.4	81.5	79.2	73.3	85.5	79.6	74.4	74.8	
HIV testing uptake, men (%)	2013	84.9	82.7	84.5	77.9	91.0	84.1	87.4	81.1	79.4	88.6	79.8	79.5	92.4	82.8	85.6	81.8	77.4	84.9	74.8	88.4	76.8	77.8	88.4	79.1	70.9	72.8	81.4	89.8	77.1	77.3	80.2	83.1	85.2	78.5	80.0	81.7	76.5	85.8	86.9	74.5	72.1	79.1	86.3	82.9	83.5	86.6	74.8	82.3
HIV testing coverage, MSM (%)	2013	86.1	81.3	90.6	81.5	90.1	86.2	89.8	89.4	84.3	80.4	86.2	90.8	90.6	88.7	84.3	81.6	93.8	89.1	80.0	81.3	90.8	89.4	79.8	85.0	85.5	76.1	88.3	90.2	76.0	87.6	91.4	87.8	90.7	83.0	84.9	88.4	91.8	85.3	79.8	94.6	86.1	82.4	85.0	83.0	85.5	88.1	94.2	91.0
HIV testing coverage, women (%)	2013	65.6	72.4	71.0	67.2	83.5	73.4	75.5	71.7	62.5	66.0	68.5	67.1	85.2	76.4	66.6	65.3	65.1	66.1	73.4	75.9	71.9	49.2	66.8	71.0	62.4	68.2	72.2	81.1	65.0	71.2	71.5	74.3	72.7	75.4	62.3	71.1	76.1	61.8	75.9	69.7	56.7	74.2	60.8	67.1	69.4	61.0	71.1	67.9
HIV testing coverage, men (%)	2013	77.5	76.9	75.3	74.3	84.0	75.8	80.4	75.0	76.0	73.1	77.9	76.9	87.1	80.1	75.1	68.7	77.5	78.2	73.3	78.9	76.0	75.0	78.1	72.5	68.4	76.4	81.2	83.3	72.9	74.3	75.6	76.8	78.2	76.7	70.5	77.8	77.7	75.6	76.7	73.7	69.6	75.4	78.3	74.3	75.4	77.7	72.9	77.6
HIV late diagnosis (%) (PHOF indicator 3.04)	2011 - 13	45.0	57.1	72.7	77.3	66.7	54.5	100.0	42.9	53.3	40.5	50.0	46.7	46.4	47.1	60.0	-	90.0	42.9	-	45.5	36.4	42.3	47.1	54.5	40.0	43.8	50.7	-	33.3	35.7	70.0	47.2	61.9	46.2	50.0	63.6	49.0	53.3	46.2	50.0	40.0	30.0	66.7	63.2	-	51.9	62.5	50.0
			< 25	25 to 50	≥ 50																																												
HIV diagnosed prevalence rate / 1,000 aged 15-59	2013	2.14	0.73	1.47	2.17	0.83	0.87	1.43	0.62	1.31	1.43	0.54	1.29	1.35	1.26	1.21	0.60	0.96	1.21	0.96	1.08	1.07	2.67	1.86	1.18	1.38	0.84	4.07	0.62	0.75	1.26	0.92	2.17	1.72	0.73	0.80	0.82	2.99	1.05	0.86	2.33	0.51	0.66	1.09	2.10	0.68	2.79	0.96	1.95

Reproductive Health

Pelvic inflammatory disease (PID) admissions rate / 100,000	2013/14	238.7	405.2	231.3	212.2	193.4	281.0	225.1	59.3	262.4	74.5	280.9	186.5	117.4	196.0	224.5	151.9	138.8	283.2	344.6	343.2	233.1	295.4	222.9	253.2	214.6	373.3	307.6	173.8	249.5	140.3	93.8	115.1	127.9	196.2	54.4	104.5	205.0	179.2	514.8	91.4	200.0	229.2	184.3	289.3	153.7	193.2	253.1	154.1
Ectopic pregnancy admissions rate / 100,000	2013/14	89.6	79.6	77.1	88.7	65.7	72.5	165.1	89.0	48.2	40.4	82.2	72.2	71.1	58.0	87.7	88.6	56.3	50.7	89.1	71.8	113.6	59.1	106.4	69.6	78.7	-	134.0	102.2	44.8	78.4	-	57.5	81.9	49.1	54.4	85.5	86.2	80.5	88.6	114.3	59.5	107.3	71.3	88.6	48.9	122.5	86.2	77.1
Cervical cancer registrations rate / 100,000	2010 - 12	9.2	9.0	7.4	8.4	8.9	10.6	8.3	8.8	5.7	8.6	7.9	5.5	5.8	6.3	10.5	8.5	3.4	8.2	14.2	12.3	11.5	10.7	5.3	6.6	8.6	11.9	10.7	12.3	7.0	8.0	11.3	12.3	11.2	6.8	4.6	9.3	7.0	4.6	5.9	8.0	10.1	8.1	7.2	5.8	5.3	10.6	12.6	5.7

Note: * - Disclosure control applied

Three Rivers

Teenage Pregnancy

Period	England	Babergh	Basildon	Bedford	Braintree	Breckland	Brentwood	Broadland	Broxbourne	Cambridge	Castle Pt	Central Beds	Chelmsford	Colchester	Dacorum	E Cambs	East Herts	Epping For	Fenland	Forest Heath	Gt Yarmouth	Harlow	Hertsmere	Hunts	Ipswich	KL & W Norf	Luton	Maldon	Mid Suffolk	North Herts	N Norfolk	Norwich	Peterboro	Rochford	S Cambs	S Norfolk	Southend	St Albans	St Edmundsbury	Stevenage	Suff Coastal	Tendring	Thr Rivers	Thurrock	Uttlesford	Watford	Waveney	Welw Hat'd	
Under 18s conception rate / 1,000 (PHOF indicator 2.04)	2013	24.3	15.4	32.9	22.0	22.3	19.3	12.8	15.5	12.5	21.9	17.4	19.9	15.8	20.6	21.0	8.6	12.2	15.4	23.1	19.9	26.5	38.3	17.0	15.4	29.4	28.7	24.4	27.1	16.3	15.5	19.0	31.4	33.4	16.8	13.4	16.0	26.6	7.6	19.1	29.1	12.8	30.3	20.2	36.1	13.3	18.5	23.3	16.7
Under 18s conceptions leading to abortion (%)	2013	51.1	40.0	51.4	50.7	55.0	55.6	89.5	54.3	77.3	42.9	44.4	58.7	57.8	44.1	56.9	-*	64.7	78.8	-*	-*	37.5	50.9	53.1	54.2	43.9	36.1	52.6	72.4	-*	52.8	46.9	42.9	46.6	63.0	59.5	54.1	37.3	-*	47.2	50.0	60.0	40.8	-*	57.4	59.1	80.0	31.9	74.2
Under 18s births rate / 1,000	2013	7.8	2.5	11.3	9.5	7.1	6.4	2.0	2.2	1.1	6.3	3.2	6.9	2.8	8.4	4.7	4.6	2.2	3.7	11.6	6.3	12.2	12.1	3.7	4.8	12.0	13.6	7.8	4.7	3.4	4.7	3.6	14.0	13.6	4.3	5.8	5.2	10.9	3.4	4.2	10.6	4.3	12.4	3.5	8.0	3.6	6.2	13.9	2.7

Wider Determinants of Health

Under 18s alcohol-specific hospital admissions rate / 100,000	2011/12 - 13/14	40.1	38.8	12.4	24.8	22.5	20.3	12.9	25.2	17.2	36.9	15.7	21.0	23.4	46.6	24.3	23.2	19.0	26.6	41.6	25.8	50.3	24.1	31.7	43.5	30.3	32.8	23.3	21.8	18.2	11.8	13.9	41.8	45.2	11.7	16.9	24.5	32.8	18.5	30.4	32.5	24.2	41.3	31.5	13.7	12.6	37.2	33.9	25.9
Percentage people living in 20% most deprived areas in England	2013	20.4	0.0	20.6	10.7	1.1	1.7	0.0	0.0	2.4	2.5	3.2	2.1	2.0	4.1	0.0	0.0	0.0	1.3	13.4	0.0	27.9	8.1	1.6	0.0	27.3	12.1	27.5	0.0	0.0	1.3	1.1	28.3	34.1	1.8	0.0	0.0	23.1	0.0	2.3	0.0	17.3	1.9	13.3	0.0	0.0	15.3	1.5	
Under 16s in poverty (%) (PHOF indicator 1.01ii)	2012	19.2	12.5	22.4	17.6	14.3	15.2	10.8	9.2	18.5	14.9	16.8	13.1	12.7	16.3	13.9	10.0	8.1	14.4	20.6	15.0	24.9	20.3	12.7	11.6	20.7	17.4	22.4	13.0	9.8	12.4	14.9	27.1	22.0	10.2	8.4	10.9	21.7	8.2	11.7	19.6	10.5	24.9	11.0	20.8	7.9	13.8	21.8	14.8
GCSE achieved 5A*-C inc. Eng & Maths (%)	2012/13	60.8	53.1	56.0	59.7	52.3	49.5	69.8	59.8	60.3	63.6	59.3	57.7	61.4	65.0	57.2	62.2	71.4	68.5	52.2	44.9	47.3	52.4	63.9	55.3	49.3	53.0	58.7	54.5	61.3	68.0	54.4	44.6	56.2	60.6	69.9	66.4	61.9	73.9	61.1	50.1	61.3	59.9	76.9	59.5	64.3	73.7	47.2	66.5
Pupil absence (%) (PHOF indicator 1.03)	2012/13	5.26	5.14	5.92	5.29	5.37	6.08	5.10	5.24	5.39	5.45	5.72	5.34	5.24	5.10	5.41	5.29	4.60	5.13	6.12	5.84	5.98	4.83	5.30	5.19	6.07	6.13	4.98	5.36	5.27	4.71	5.88	6.04	5.39	5.11	5.22	4.92	5.36	4.49	5.17	5.71	5.30	5.77	4.78	5.23	4.94	5.35	6.10	4.86

Comparison with respect to England value / goal




Note: * - Disclosure control applied


Bar Charts

Key Indicators


Syphilis diagnosis rate / 100,000




Gonorrhoea diagnosis rate / 100,000




Chlamydia detection rate / 100,000 aged 15-24 (PHOF indicator 3.02)




Chlamydia proportion aged 15-24 screened




All new STI diagnoses (exc Chlamydia aged <25) / 100,000




HIV testing uptake, MSM (%)




HIV late diagnosis (%) (PHOF indicator 3.04)




HIV diagnosed prevalence rate / 1,000 aged 15-59



Under 18s conception rate / 1,000 (PHOF indicator 2.04)



Under 18s conceptions leading to abortion (%)




Key --- England value and confidence interval ↑ Three Rivers Other local authority in East of England

Three Rivers

Key Indicators continued


Sexual offences rate / 1,000 (PHOF indicator 1.12iii)




Key  England value and confidence interval  Three Rivers  Other local authority in East of England

HIV & STI


Syphilis diagnosis rate / 100,000




Gonorrhoea diagnosis rate / 100,000




Chlamydia detection rate / 100,000 aged 15-24 (PHOF indicator 3.02)




Chlamydia proportion aged 15-24 screened




Genital warts diagnosis rate / 100,000




Genital herpes diagnosis rate / 100,000




All new STI diagnoses (exc Chlamydia aged <25) / 100,000




STI testing rate (exc Chlamydia aged < 25) / 100,000



STI testing positivity (exc Chlamydia aged <25) %



HIV testing uptake, MSM (%)




Key England value and confidence interval ↑ Three Rivers Other local authority in East of England


Three Rivers

HIV & STI continued


HIV testing uptake, women (%)




HIV testing uptake, men (%)




HIV testing coverage, MSM (%)




HIV testing coverage, women (%)




HIV testing coverage, men (%)



HIV late diagnosis (%) (PHOF indicator 3.04)




HIV diagnosed prevalence rate / 1,000 aged 15-59




Key England value and confidence interval ↑ Three Rivers Other local authority in East of England

Reproductive Health


Pelvic inflammatory disease (PID) admissions rate / 100,000



Ectopic pregnancy admissions rate / 100,000



Cervical cancer registrations rate / 100,000




Key England value and confidence interval Three Rivers Other local authority in East of England


Three Rivers

Teenage Pregnancy


Under 18s conception rate / 1,000 (PHOF indicator 2.04)



Under 18s conceptions leading to abortion (%)




Under 18s births rate / 1,000




Key England value and confidence interval ↑ Three Rivers Other local authority in East of England

Wider Determinants of Health


Under 18s alcohol-specific hospital admissions rate / 100,000




Percentage people living in 20% most deprived areas in England




Under 16s in poverty (%) (PHOF indicator 1.01ii)



GCSE achieved 5A*-C inc. Eng & Maths (%)



Pupil absence (%) (PHOF indicator 1.03)



Key England value and confidence interval ↑ Three Rivers Other local authority in East of England

Three Rivers

Definitions

Key Indicators

- Syphilis diagnosis rate / 100,000** - All syphilis diagnoses among people accessing GUM clinics in England who are also residents in England, expressed as a rate per 100,000 population. Data excludes people accessing GUM clinics located in England who are residents in Wales, Scotland, Northern Ireland or abroad.
- Gonorrhoea diagnosis rate / 100,000** - All gonorrhoea diagnoses among people accessing GUM clinics in England who are also residents in England, expressed as a rate per 100,000 population. Data excludes people accessing GUM clinics located in England who are residents in Wales, Scotland, Northern Ireland or abroad.
- Chlamydia detection rate / 100,000 aged 15-24 (PHOF indicator 3.02)** - All chlamydia diagnoses among 15 to 24 year olds accessing GUM clinics, primary care and community services in England who are also residents in England, expressed as a rate per 100,000 population. Data excludes people accessing services located in England who are residents in Wales, Scotland, Northern Ireland, abroad or where an area of residence cannot be attributed.
- Chlamydia proportion aged 15-24 screened** - All chlamydia tests (asymptomatic screens and symptomatic tests) undertaken in 15 to 24 year olds attending GUM clinics, primary care and community services who are residents in England. Data excludes people accessing services located in England who are residents in Wales, Scotland, Northern Ireland or abroad.
- All new STI diagnoses (exc Chlamydia aged <25) / 100,000** - All new STI diagnoses (excluding Chlamydia in under 25 year olds) in GUM clinic attendees who are resident in England, i.e. data excludes people accessing GUM clinics located in England who are resident in Wales, Scotland, Northern Ireland or abroad.
- HIV testing uptake, MSM (%)** - The number of 'Eligible new GUM Episodes' where a HIV test was accepted as a proportion of those where a HIV test was offered. Multiple episodes of HIV test offer and uptake are included per individual within a year. 'Eligible new GUM Episode' is defined as a visit to a GUM clinic including all subsequent GUM attendances in the following six weeks (i.e. eligibility for testing occurs only once every six weeks). Attendances by known HIV positive patients¹, or where a HIV test was not appropriate² are excluded. ¹ E1B, E2B, E3A2, E3B, H, H2 ² P1C Data excludes people accessing GUM clinics located in England who are residents in Wales, Scotland, Northern Ireland or abroad. MSM includes men who reported a homosexual or bisexual orientation. For uptake of HIV testing, MSM is defined based on a patient's entire clinic attendance history (i.e. a man is classified as MSM for all attendance years including and following the earliest year a man identifies as MSM).
- HIV late diagnosis (%) (PHOF indicator 3.04)** - All adults (aged 15+ years) newly diagnosed with HIV infection with CD4 counts available within 91 days of diagnosis and with known residence based information.
- HIV diagnosed prevalence rate / 1,000 aged 15-59** - People aged 15 to 59 years who were seen at HIV care services.
- Under 18s conception rate / 1,000 (PHOF indicator 2.04)** - Conceptions in women aged under 18 per 1,000 females aged 15-17
- Under 18s conceptions leading to abortion (%)** - The percentage of conceptions to those aged under 18 years that led to an abortion
- Sexual offences rate / 1,000 (PHOF indicator 1.12iii)** - Rate of sexual offences based on police recorded crime data per 1,000 population

HIV & STI

- Syphilis diagnosis rate / 100,000** - All syphilis diagnoses among people accessing GUM clinics in England who are also residents in England, expressed as a rate per 100,000 population. Data excludes people accessing GUM clinics located in England who are residents in Wales, Scotland, Northern Ireland or abroad.
- Gonorrhoea diagnosis rate / 100,000** - All gonorrhoea diagnoses among people accessing GUM clinics in England who are also residents in England, expressed as a rate per 100,000 population. Data excludes people accessing GUM clinics located in England who are residents in Wales, Scotland, Northern Ireland or abroad.
- Chlamydia detection rate / 100,000 aged 15-24 (PHOF indicator 3.02)** - All chlamydia diagnoses among 15 to 24 year olds accessing GUM clinics, primary care and community services in England who are also residents in England, expressed as a rate per 100,000 population. Data excludes people accessing services located in England who are residents in Wales, Scotland, Northern Ireland, abroad or where an area of residence cannot be attributed.
- Chlamydia proportion aged 15-24 screened** - All chlamydia tests (asymptomatic screens and symptomatic tests) undertaken in 15 to 24 year olds attending GUM clinics, primary care and community services who are residents in England. Data excludes people accessing services located in England who are residents in Wales, Scotland, Northern Ireland or abroad.
- Genital warts diagnosis rate / 100,000** - All diagnoses of first episode genital warts among people accessing GUM clinics in England who are also residents in England, expressed as a rate per 100,000 population. Data excludes people accessing GUM clinics located in England who are resident in Wales, Scotland, Northern Ireland or abroad.
- Genital herpes diagnosis rate / 100,000** - All diagnoses of first episode genital herpes among people accessing GUM clinics in England who are also residents in England, expressed as a rate per 100,000 population. Data excludes people accessing GUM clinics located in England who are resident in Wales, Scotland, Northern Ireland or abroad.
- All new STI diagnoses (exc Chlamydia aged <25) / 100,000** - All new STI diagnoses (excluding Chlamydia in under 25 year olds) in GUM clinic attendees who are resident in England, i.e. data excludes people accessing GUM clinics located in England who are resident in Wales, Scotland, Northern Ireland or abroad.
- STI testing rate (exc Chlamydia aged <25)** - All tests for syphilis, HIV, *Neisseria gonorrhoeae* and chlamydia (aged over 25) in GUM clinic attendees who are resident in England, i.e. data excludes people accessing GUM clinics located in England who are resident in Wales, Scotland, Northern Ireland or abroad.
- STI testing positivity (exc Chlamydia aged <25) %** - All new STI diagnoses (excluding Chlamydia in under 25 year olds) as a percentage of the number of STI tests performed.
- HIV testing uptake, MSM (%)** - The number of 'Eligible new GUM Episodes' where a HIV test was accepted as a proportion of those where a HIV test was offered. Multiple episodes of HIV test offer and uptake are included per individual within a year. 'Eligible new GUM Episode' is defined as a visit to a GUM clinic including all subsequent GUM attendances in the following six weeks (i.e. eligibility for testing occurs only once every six weeks). Attendances by known HIV positive patients¹, or where a HIV test was not appropriate² are excluded. ¹ E1B, E2B, E3A2, E3B, H, H2 ² P1C Data excludes people accessing GUM clinics located in England who are residents in Wales, Scotland, Northern Ireland or abroad. MSM includes men who reported a homosexual or bisexual orientation. For uptake of HIV testing, MSM is defined based on a patient's entire clinic attendance history (i.e. a man is classified as MSM for all attendance years including and following the earliest year a man identifies as MSM).

HIV testing coverage, MSM (%) - The proportion of 'Eligible new GUM Attendees' in whom a HIV test was accepted. 'Eligible new GUM Attendee' is defined as a patient attending a GUM clinic at least once during a calendar year. Patients known to be HIV positive¹, or for whom a HIV test was not appropriate² are excluded ¹ E1B, E2B, E3A2, E3B, H, H2 ² P1C Data excludes people accessing GUM clinics located in England who are residents in Wales, Scotland, Northern Ireland or abroad. MSM includes men who reported a homosexual or bisexual orientation. For coverage of HIV testing, MSM is defined based on a patient's entire clinic attendance history (i.e. a man is classified as MSM for all attendance years including and following the earliest year a man identifies as MSM).

HIV late diagnosis (%) (PHOF indicator 3.04) - All adults (aged 15+ years) newly diagnosed with HIV infection with CD4 counts available within 91 days of diagnosis and with known residence based information.

HIV diagnosed prevalence rate / 1,000 aged 15-59 - People aged 15 to 59 years who were seen at HIV care services.

Reproductive Health

Pelvic inflammatory disease (PID) admissions rate / 100,000 - Crude rate of pelvic inflammatory disease (PID) admissions in women aged 15-44 years per 100,000 population (women aged 15-44 years)

Ectopic pregnancy admissions rate / 100,000 - Crude rate of ectopic pregnancy admissions to hospital in women aged 15-44 years per 100,000 population (women aged 15-44 years)

Cervical cancer registrations rate / 100,000 - Directly age-standardised rate of cervical cancer registrations per 100,000 female all age population

Teenage Pregnancy

Under 18s conception rate / 1,000 (PHOF indicator 2.04) - Conceptions in women aged under 18 per 1,000 females aged 15-17

Under 18s conceptions leading to abortion (%) - The percentage of conceptions to those aged under 18 years that led to an abortion

Under 18s births rate / 1,000 - Births in women aged under 18 per 1,000 females aged 15-17

Wider Determinants of Health

Under 18s alcohol-specific hospital admissions rate / 100,000 - Persons admitted to hospital due to alcohol-specific conditions – under 18 year olds, crude rate per 100,000 population. See LAPE user guide for further details - http://www.lape.org.uk/downloads/Lape_guidance_and_methods.pdf

Percentage people living in 20% most deprived areas in England - Percentage of the relevant population in this area living in the 20% most deprived Lower Super Output Areas in England.

Under 16s in poverty (%) (PHOF indicator 1.01ii) - Percentage of children in low income families (children living in families in receipt of out of work benefits or tax credits where their reported income is less than 60% median income) for under 16s only

GCSE achieved 5A*-C inc. Eng & Maths (%) - Percentage of pupils achieving 5 or more GCSEs at grades A*-C (including English and Maths) or equivalent, percentage of pupils at end of Key Stage 4 in schools maintained by the Local Authority, at the end of the academic year, persons.

Pupil absence (%) (PHOF indicator 1.03) - Percentage of half days missed by pupils due to overall absence (including authorised and unauthorised absence).